

GEORGIA INSTITUTE OF TECHNOLOGY  
OFFICE OF CONTRACT ADMINISTRATION  
SPONSORED PROJECT INITIATION

Date: August 26, 1980

Project Title: New Methods for Biomedical Research

Project No: G-33-D05

Project Director: Dr. Nai-Teng Yu

Sponsor: DHEW/PHS/NIH National Eye Institute (Research Career Development Program)

Agreement Period: From 1 July 1980 Until 30 June 1981 (05 year)

Type Agreement: Grant No. 5-K04-EY00073-05

Amount: \$35,805 G-33-D05

Reports Required: Annual Progress Reports with continuation applications; Terminal Progress Report upon Grant expiration.

Sponsor Contact Person (s):

Technical Matters

George M. Steinberg, Ph.D.  
Program Official  
Extramural Program Director  
Cataract Program  
National Eye Institute  
Bethesda, Maryland 20014

Contractual Matters

(thru OCA)

Gaye Lynch  
Grants Management Contract  
Extramural Services Branch  
National Eye Institute  
Bethesda, Maryland 20014

Phone: 301/496-5884

NOTE: FOLLOW-ON PROJECT TO G-33-D04  
(04 year)

Grants Management Officer;  
Anna Marie Perrell

Defense Priority Rating: None

Assigned to: Chemistry (School/Laboratory)

COPIES TO:

Project Director  
Division Chief (EES)  
School/Laboratory Director  
Dean/Director-EES  
Accounting Office  
Procurement Office  
Security Coordinator (OCA)

Library, Technical Reports Section  
EES Information Office  
EES Reports & Procedures  
Project File (OCA)  
Project Code (GTRI)  
Other \_\_\_\_\_

✓ Reports Coordinator (OCA)

Research Property Coordinator (OCA)

SPONSORED PROJECT TERMINATION SHEETDate 12/30/81

Project Title: New Methods for Biomedical Research

Project No: G-33-D05

Project Director: Dr. Nai-Teng Yu

Sponsor: DHEW/PHS/NIH National Eye Institute (Research Career Development Program)

Effective Termination Date: 6/30/81Clearance of Accounting Charges: 6/30/81

Grant/Contract Closeout Actions Remaining:

- ☐ Final Invoice and Closing Documents
- ☐ Final Fiscal Report
- ☒ Final Report of Inventions
- ☐ Govt. Property Inventory & Related Certificate
- ☐ Classified Material Certificate
- ☐ Other \_\_\_\_\_

Assigned to: Chemistry (School/Laboratory)COPIES TO:

Administrative Coordinator  
Research Property Management  
Accounting  
Procurement/EES Supply Services

Research Security Services  
Reports Coordinator (OCA) ✓  
Legal Services (OCA)  
Library

EES Public Relations (2)  
Computer Input  
Project File  
Other \_\_\_\_\_

Final Progress Report

Grant Number: 5K04EY00073  
Period Covered: 7/1/76 - 6/30/81  
Investigator: Yu, Nai-Teng  
Institution: Georgia Institute of Technology  
Atlanta, GA 30332  
Title of Project: New Methods for Biomedical Research

The most important accomplishment during the period supported by the N.I.H. Research Career Development Award is the development of Raman scattering technique to a stage that one can obtain high quality Raman spectra of the lens from a live animal without injury to the eye. Raman spectroscopy has been shown to be a useful tool in the study of lens structure (Yu and East, 1975; Kuck, East and Yu, 1976; Yu, East, Chang and Kuck, 1977; East, Chang, Yu and Kuck, 1978; Kuck, Yu and Askren, 1981). Recent applications of the Raman probe have led to the discovery of different fluorescent materials in the lens (Kuck and Yu, 1978; Yu, Kuck and Askren, 1979). Of particular interest is one fluorophor with excitation/emission maxima: 647/672 nm (Kuck, Yu and Askren, 1979). This fluorophor presents itself as a precatactous indicator since it does not appear in the normal young lens, but is highly elevated in brunescant cataracts.

We recognized the possible clinical application of our technique and instrumentation, with some refinement. The ultimate goal of this research is the clinical application as a routine diagnostic procedure in ophthalmology.

In a step-wise fashion we proceeded towards obtaining Raman spectra from live animals. Isolated eyes from rabbits were used first in order to work out orientation and alignment details. An arrangement with a 90° scattering angle was chosen for simplicity. Raman spectra from the rabbit lens in situ can be obtained with a 12 mW laser power at 514.5 nm wavelength.

Human eyes (64 yr. old) were tried, but fluorescence interfered with data collection. With modification, the Raman instrumentation can be used for measuring the fluorophor concentration. Only human eyes with the cornea intact were used. Since most corneas are removed for transplant surgery, these eyes were rare and we had the opportunity of working with them only once.

Pentobarbital was used as an anesthetic initially; later, tholazine was added to the protocol as a "pre-op" medication. Rabbits, even under anesthesia, have a propensity towards twitching. Sneezes, pulsations and other irritations would make the eye move out of alignment. This made spectrum-scanning with the conventional single-channel photo-multiplier tube impossible. Spectrum scanning from 200-3600  $\text{cm}^{-1}$  took thirty minutes to run and the animal had to be motionless throughout. This problem was remedied with the installation of a Reticon detector (see Fig. 1). This instrument has a multiple photodiode arrays whose

different pixels accumulate signal simultaneously. The data is integrated over time and therefore inconsistencies in the sample scattering are inconsequential. The position of the eye in the laser beam could be quickly realigned with little change in the quality of the spectrum obtained.

A further advantage of this detection system is that a reduced laser power could be used. Fig. 2 shows the data of Mathies and Yu (1978) who compared the quality of spectra obtained from an intact bovine lens using a conventional photomultiplier and a Silicon Intensified Target. They were able to obtain satisfactory spectra with as little as 1 mW laser power using the SIT. This detector, however, does not have the sensitivity of the Reticon we have used in the laser wavelength region. The increased sensitivity of our detector has permitted the acquisition of suitable spectra in as little as 15 seconds, with excellent spectra in 2-1/2 minutes, whereas the spectra of Mathies and Yu required 15 minutes or more.

The only disadvantage of our set-up is that it is designed for high resolution and therefore it has a greater dispersion. Therefore, several "snap-shorts" of the spectrum must be made in order to cover the complete spectrum. Each spectrum consists of 5 pieces that were spliced together.

### Conclusions

The structural information obtainable from the Raman spectrum has been documented, but nowhere in the literature has a spectrum from the lens of a live organism been reported. This accomplishment represents a large step towards the use of this technique in the physician's office.

The difficulties that has yet to be overcome include the dispersion and final design. With a monochromator design for a laser degree of dispersion, only a single "exposure" will be necessary to obtain a spectrum. In the interest of obtaining a fluorescent spectrum from a brunescant cataract, an even shorter exposure time than that needed for Raman spectroscopy will provide a suitable spectrum. This is in part due to the fact that fluorescence is generally more intense than Raman scattering but also due to the "squeezing" effect of the decreased dispersion. Also, fluorescent spectroscopy does not require high resolution.

With increased use of lasers in Medicine, particularly in the field of ophthalmology, only minor modification of current diagnostic and therapeutic instrumentation need to be made before this technique will be ready for clinical investigation. Information about the clinical development (etiology) of brunescant cataracts that this technique could provide will be useful to the ophthalmologist.

### References:

Askren, C. C., Yu, N. T. and Kuck, J. F. R. (1979) Exp. Eye Res. 29, 647.

- East, E. J., Chang, R. C. C., Yu, N. T. and Kuck, J. R. F. (1978) J. Biol. Chem. 253, 1436.
- Kuck, J. F. R., East, E. J. and Yu, N. T. (1976) Exp. Eye Res. 23, 9.
- Kuck, J. F. R. and Yu, N. T. (1978) Exp. Eye Res. 27, 737.
- Kuck, J. F. R., Yu, N. T. and Askren, C. C. (1981) Exp. Eye Res. (in press).
- Mathies, R. and Yu, N. T. (1978) J. Raman Spectros. 7, 349.
- Yu, N. T. and East, E. J. (1975) J. Biol. Chem., 250, 2195.
- Yu, N. T., East, E. J., Chang, R. C. C. and Kuck, J. F. R. (1977) Exp. Eye Res. 24, 321.
- Yu, N. T., Kuck, J. F. R. and Askren, C. C. (1979) Invest. Ophthalm. and Vis. Sci. 18, 1278.

Publications which acknowledge the support by EY00073 Award:

1. Yu, N. T., East, E. J., Chang, R. C. and Kuck, J. F. R., Jr. (1977) "Raman Spectra of Bird and Reptile Lens Proteins" Exp. Eye Res. 24, 321.
2. Yu, N. T. (1977) "Raman Spectroscopy" A Conformational Probe in Biochemistry" CRC Critical Reviews in Biochemistry, 4, 229.
3. Shelnut, J. A., Cheung, L. D., Chang, R. C. C., Yu, N. T. and Felton, R. H. (1977) "Resonance Raman Spectra of Metalloporphyrins. Effects of Jahn-Teller Instability and Nuclear Distortion on Excitation Profiles of Stokes Fundamental" J. Chem. Phys. 66, 3387.
4. Chang, L. D., Yu, Nai-Teng and Felton, R. H. (1978) "Resonance Raman Spectra and Excitation Profiles of Soret-Excited Metalloporphyrins" Chem. Phys. Lett., 55, 527.
5. Felton, R. H. and Yu, Nai-Teng (1978) "Resonance Raman Scattering from Metalloporphyrins and Hemoproteins" in The Porphyrins (Dolpnin, D., Ed.) Vol. III, Chapter VIII, pp. 347-343, Academic Press, New York.
6. Mathies, R. and Yu, N. T. (1978) "Raman Spectroscopy with Intensified Vidicon Detectors: A Study of Intact Bovine Lens Proteins" J. Raman Spectrosc., 7, 349.
7. Kuck, J. F. R. and Yu, N. T. (1978) "Raman and Fluorescent Emission of the Human Lens. A New Fluorophor" Exp. Eye Res. 27, 737.
8. Yu, N. T. and Kuck, J. F. R. (1978) "Focusing on Lenses with Laser Raman Spectroscopy" The Spex Speaker, Vol. 23, No. 3, pp. 1-8 (Published by Spex Industries, Inc., 3380 Park Ave., Metchen, NJ. 08840).



9. East, E. J., Chang, R. C. C., Yu, N. T. and Kuck, J. F. R. (1978) "Raman Spectroscopic Measurement of Total Sulfhydryl in Intact Lens as Affected by Aging and Ultraviolet Irradiation. Deuterium Exchange as a Probe for Accessible Sulfhydryl in Living Tissue", J. Biol. Chem. 253 1436.
10. Yu, N. T., Kuck, J. F. R. and Askren, C. C. (1979) "Red Fluorescence in Older and Brunescent Human Lenses" Invest. Ophthal. & Vis. Sci., 18, 1278.
11. Lanir, A. and Yu, N. T. (1979) "A Raman Spectroscopic Study of the Interaction of Divalent Metal Ions with Adenine Moiety of Adenosine 5'-Tri-phosphate" J. Biol. Chem. 254, 5882.
12. Askren, C. C., Yu, N. T. and Kuck, J. F. R. (1979) Variation of the Concentration of Sulfhydryl along the Visual Axis of Aging Lenses by Laser Raman Optical Dissection Technique" Exp. Eye Res. 29, 647.
13. Lanir, A., Yu, N. T. and Felton, R. H. (1979) "Conformational Transitions and Vibronic Couplings in Acid Ferricytochrome c: A Resonance Raman Study", Biochemistry, 18, 1656.
14. Yu, N. T. and Srivastava, R. B. (1980) "Resonance Raman Spectroscopy of Heme Proteins with Intensified Vidicon Detectors: Studies of Low Frequency Modes and Excitation Profiles in Cytochrome c and Hemoglobin" J. Raman Spectrosc., 9, 166.
15. Yu, N. T. and Tsubaki, M. (1980) "Resonance Raman Spectra of Manganese Myoglobin and Its Azide Complex. Assignment of a New Charge-Transfer Band to Azide ( $\pi$ )  $\rightarrow$  Porphyrin ( $\pi^*$ ) Transition" Biochemistry, 19, 4647.
16. Tsubaki, M., Srivastava, R. B. and Yu, N. T. (1981) "Temperature Dependence of Resonance Raman Spectra of Metmyoglobin and Methemoglobin Azide. Detection of Resonance-Enhanced Bound Azide Vibrations and Iron-Azide Stretch" Biochemistry, 20, 946.
17. Srivastava, R. B., Pace, C. and Yu, N. T. (1981) "Comparative Raman Studies of Cytochrome b<sub>562</sub> and Cytochrome c" J. Raman Spectrosc. 11, 20.
18. Yu, N. T., Lanir, A. and Werber, M. M. (1981) "Laser Raman Scattering and Resonance in Co(III)-ATP Complexes", J. Raman Spectrosc. 11, 150 (1981).
19. Tsubaki, M. and Yu, N. T. (1981) "Resonance Raman Investigation of Dioxygen Bonding in Oxycobalmyoglobin and Oxycobalthemoglobin. Structural Implication of Splittings of Bound O-O Stretching Vibration" Proc. Natl. Acad. Sci., USA 78, 3581.
20. Tsubaki, M., Srivastava, R. B. and Yu, N. T. (1981) "Resonance Raman Investigation of Carbon Monoxide Bonding in Carbonmonoxy Hemoglobin and Myoglobin: Detection of Fe-CO Stretching and Fe-C-O Bending Vibrations and Influence of the Quaternary Structure Change" Biochemistry (in press).

21. Tsubaki, M. and Yu, N. T. (1981) "Resonance Raman Investigation of Nitric Oxide Bonding and Nitrosyl Hemoglobin A and Myoglobin" Detection of Bound N-O Stretch and Fe-NO Stretch Vibrations of Hexa-coordinated NO-Heme Complex" Biochemistry (in press).
  22. Kuck, J. F. R., Yu, N. T. and Askren, C. C. (1981) "Total Sulfhydryl by Raman Spectroscopy in the Intact Lens of Several Studies: Variations in the Nucleus and Along the Optical Axis During Aging" Exp. Eye Res. (in press).
  23. Yu, N. T., Mackin, H. C. and Kerr, E. A. (1982) "Raman Spectroscopy of Heterocyclic Compounds" in Physical Methods in Heterocyclic Chemistry (Gupta, R. R., Ed.) John-Wiley Interscience Publishers, New York.
  24. Mackin, H. C., Tsubaki, M. and Yu, N. T. (1982) "Resonance Raman Spectra of CoO<sub>2</sub> Moiety in Oxy Co "Picket Fence" Porphyrin and Oxy Co Mesoporphyrin IX-substituted Myoglobin. Resonance Interaction between O-O Stretch and Porphyrin Ring Vibrations" J. Am. Chem. Soc. (in press).
- 
25. Kerr, E., Mackin, H. C. and Yu, N. T. (1982) "Influence of Strong or Weak Field Trans Ligands upon the Stretch of Iron-Carbon Bond in Carbonmonoxy Iron "Picket Fence" Porphyrin as Detected by Resonance Raman Spectroscopy" (in preparation).






Fig. 1





Raman spectra of low frequency vibrations of bovine lens proteins using PMT-PC detection. (A) Spectrum taken with 100 mW 514.5 nm and  $5 \text{ cm}^{-1}$  ( $300 \mu$ ) slits. Data acquisition time 800 sec. (B) Four-fold expansion of A. (C) Spectrum taken with 1 mW 514.5 nm and  $6.7 \text{ cm}^{-1}$  ( $400 \mu$ ) slits. Data acquisition time 800 sec.



Raman spectra of low frequency vibrations of bovine lens proteins using SIT-OMA detection. (A) Spectrum taken with 100 mW 514.5 nm, no delays, and  $10.5 \text{ cm}^{-1}$  ( $150 \mu$ ) slits. Total data integration time 29.5 sec (900 cycles). One bin is equal to  $1.54 \text{ cm}^{-1}$ . (B) Four-fold expansion of A. (C) Spectrum taken with 1 mW 514.5 nm,  $14 \text{ cm}^{-1}$  ( $200 \mu$ ) slits, and 100 delay cycles. Total data integration time 655 sec (20,000 cycles).

Fig. 2




Fig. 3